



## **You had Partial Knee Replacement with Dr. Brannen**

- **Follow-up** with Dr. Brannen's office at your 2-week post-operative visit
- **Activity:** You are weight bearing as tolerated with a Walker
  - ✓ Keep pressure ice pack on for at least 30 minutes on then 1 hour off. Repeat this as necessary. Never put ice directly on skin.
  - ✓ Use your Incentive Spirometer (inhale) 10 x an hour (while awake) for 14 days. Deep breathe in through your nose, hold it for 3 seconds and exhale.
- **Medications:** Take your pain medicine exactly as directed. **Do not drive if you have taken any prescription pain medication.** Pain medication can lead to constipation. High fiber diet, lots of fluids, and a stool softener (Colace) or laxative (Miralax)
  - ☐ **Norco (Hydrocodone/acetaminophen)** 1-2 tabs every 4-6 hours as needed for pain
  - ☐ **Aspirin 325 mg** 1 tab once a day for 30 days – First dose day of surgery
  - ☐ **Ibuprofen 600mg** every 6 hrs a day - First dose tomorrow
  - ☐ **Cefadroxil 500 mg 1** tab twice a day – First dose tomorrow if you had an outpatient procedure
- **Blood Clot Prevention:**
  - Perform ankle pump exercises (moving your foot back and forth) for a minimum 14 days.
- **Surgical Dressing:**
  - You may shower with the dressing covered
  - Wrap in plastic saran wrap, or Press and Seal, when showering then remove wrapping.
  - If the bandage gets wet. Please remove. Do not use other tape to hold it on.
  - No hot tubs, pools, or baths until cleared by your healthcare provider.